

Thirteen Things Physicians and Patients Should Question

1 Don't use atypical antipsychotics as a first-line intervention for insomnia in children and youth.

Recent research confirms a dramatic increase in the use of atypical antipsychotics with subsequent side-effects including obesity, which is already a major health issue. It is prudent to pursue nonpharmacological measures first, such as behavioural modifications and ensuring good sleep hygiene (such as eliminating daytime napping and shutting off electronics an hour before bedtime). If these interventions are not successful, then consider short-term use of melatonin.

2 Don't use SSRIs as the first-line intervention for mild to moderately depressed teens.

Evidence clearly indicates that antidepressant medication is less effective in children and adolescents up to the age of 17 years and first-line treatment for this group should include cognitive behavioural therapy or interpersonal psychotherapy. Attention should always be focused on children's and teens' environmental safety and adequate parental support to avoid missing cases of neglect or abuse. Following this, a first-line intervention should be psychoeducation on the importance of regular sleep, diet and exercise to ensure healthy, age-appropriate developmental support.

3 Don't use atypical antipsychotics as a first-line intervention for Attention Deficit Hyperactivity Disorder (ADHD) with disruptive behaviour disorders.

Treatment of ADHD should include adequate education of patients and their families, behavioural interventions, psychological treatments and educational accommodations first. If this approach is not sufficient, stimulant medication and a behavioural analysis to ensure appropriate support from the parent and classroom is indicated. The use of alpha 2 agonists (such as guanfacine) and atomoxetine should be considered before using atypical antipsychotics (such as risperidone) in children with disruptive behaviour disorders (oppositional defiant disorder, conduct disorder).

4 Don't use psychostimulants as a first-line intervention in preschool children with ADHD.

Preschool children with ADHD need to be assessed for other neurodevelopmental disorders and consideration given to environmental stressors such as neglect, abuse or exposure to domestic violence. Treatment also includes adequate education and support of parents followed by advice on behavioural management and community placement.

5 Don't routinely use antipsychotics to treat primary insomnia in any age group.

Second-generation antipsychotics (SGAPs), such as olanzapine and quetiapine, have sedative properties, and are often prescribed off-label for complaints of insomnia. These drugs carry significant risk of potential side-effects including weight gain and metabolic complications, even at low doses used to treat insomnia. In patients with dementia, they can also potentially cause serious side-effects of increased risk of cerebrovascular event and increased risk of death.

6 Don't routinely order qualitative toxicology (urine drug screen) testing on all psychiatric patients presenting to emergency rooms.

Qualitative urine toxicology testing has not been shown to improve the routine management of psychiatric patients in emergency rooms because of the potential for false positives, false negatives, true positives which are unrelated or minimally relevant to the clinical presentation, and finally the delay in psychiatric assessment and management as a result of testing.

7 Don't routinely use antidepressants as first-line treatment for mild or subsyndromal depressive symptoms in adults.

Antidepressant response rates are higher for depression of a moderate to severe nature. For mild or subsyndromal depressive symptoms a complete assessment, ongoing support and monitoring, psychosocial interventions and lifestyle modifications should be the first lines of treatment. This may avoid the side-effects of medication and establish etiological factors important to future assessment and management. Antidepressants are appropriate in cases of persistent mild depression, where there is a past history of more severe depression, or where other interventions have failed.

8 Don't routinely order brain neuroimaging (CT or MRI) in first episode psychoses in the absence of signs or symptoms suggestive of intracranial pathology.

Signs and symptoms suggestive of intracranial pathology include headaches, nausea and vomiting, seizure-like activity, and later-age of onset of symptoms. Multiple studies have found that routine neuroimaging in first episode psychoses does not yield findings which alter clinical management in a meaningful way. The risks of radiation exposure and delay in treatment also argue against routine neuroimaging.

9 Don't routinely continue benzodiazepines initiated during an acute care hospital admission without a careful review and plan of tapering and discontinuing, ideally prior to hospital discharge.

Benzodiazepines, while helpful for short-term relief of anxiety and insomnia, are associated with a variety of side-effects and long-term problems including cognitive and psychomotor impairment as well as abuse and dependence. Benzodiazepines are commonly used in hospital to treat anxiety or insomnia in association with either the presenting condition or the hospital environment. Once the presenting condition is treated, benzodiazepines should be tapered and discontinued. For patients who are still on benzodiazepines at the time of discharge, a plan for tapering and discontinuing them after discharge should be completed and specified in the discharge summary and prescription.

10 Don't routinely prescribe antidepressants as first-line treatment for depression comorbid with an active alcohol use disorder without first considering the possibility of a period of sobriety and subsequent reassessment for the persistence of depressive symptoms.

The concurrent management of psychiatric illness and alcohol use disorders requires evaluation of the role alcohol plays as a causative factor for depressive symptoms. Studies have found that response rates to antidepressants are higher when antidepressants are reserved for persistence of symptoms after a period of sobriety lasting from two to four weeks. Additionally, studies have demonstrated remission from depressive symptoms with sobriety in the absence of antidepressant treatment in a significant percentage of cases. Management of comorbid psychiatric illness and substance use disorders including alcohol dependence involves assessment and treatment delivered in a concurrent manner.

11 Don't routinely prescribe high-dose or combination antipsychotic treatment strategies in the treatment of schizophrenia.

High-dose and combination strategies involving atypical antipsychotics (AAPs) are used in clinical practice for patients with schizophrenia who are inadequately controlled with one or more AAPs used at standard doses. A recent meta-analysis found no clinically significant improvements in patients with schizophrenia who were inadequately controlled on standard-dose antipsychotics when treated with combination or high-dose AAPs. In terms of safety, no clinically significant differences were evident between combination or high-dose therapy in comparison with standard-dose monotherapy.

12 Don't use antipsychotics as first choice to treat behavioural and psychological symptoms of dementia.

People with dementia often exhibit challenging behavioural symptoms such as aggression and psychosis. In such instances, antipsychotic medicines may be necessary, but should be prescribed cautiously as they provide limited benefit and can cause serious harm, including premature death. Use of these drugs should be limited in dementia to cases where nonpharmacologic measures have failed, and where the symptoms either cause significant suffering, distress, and/or pose an imminent threat to the patient or others. A thorough assessment that includes identifying and addressing causes of behaviour change can make use of these medications unnecessary. Epidemiological studies suggest that typical (i.e., first generation) antipsychotics (i.e., haloperidol) are associated with at least the same risk of adverse events. This recommendation does not apply to the treatment of delirium or major mental illnesses such as mood disorders or schizophrenia.

13 Don't use benzodiazepines or other sedative-hypnotics in older adults as first choice for insomnia.

Nonpharmacological interventions such as cognitive behavioural therapy and brief behavioural interventions have proven benefit in the management of insomnia in older adults. Epidemiological studies have shown that the risk of motor vehicle accidents, falls and hip fractures leading to hospitalization and death can more than double in older adults taking benzodiazepines and other sedative-hypnotics. Prescribing or discontinuing sedative-hypnotics in hospital can have substantial impact on long-term use. These potential harms and others such as impaired cognition need to be recognized when considering treatment strategies for insomnia. Use of benzodiazepines should be limited to as short a period as possible, in cases where nonpharmacological therapies have failed, and the symptoms of sleep disturbance cause significant suffering or distress.

How the list was created

The Canadian Psychiatric Association (CPA) determined its *Choosing Wisely Canada* recommendations by establishing a working group that included representatives from the CPA's Professional Standards and Practice Committee, Research Committee, and Member-in-Training Section, as well as the Canadian Academy of Geriatric Psychiatry (CAGP) and the Canadian Academy of Child and Adolescent Psychiatry (CACAP). A person with lived experience from the Canadian Mental Health Association was also a member of the working group. CPA members were invited to provide suggestions for potential list items, as were the provincial psychiatric associations, the Canadian Academy of Psychiatry and the Law (CAPL) and the Canadian Academy of Psychosomatic Medicine (CAPM). The working group considered suggestions received, and assistance was obtained from the Addiction and Mental Health Strategic Clinical Network for Alberta Health Services in conducting rapid literature reviews on a number of potential CPA list items. List items were further refined in subsequent working group teleconferences, and a next-to-final draft was recirculated to the provincial psychiatric associations, CAPL and CAPM for final comments, which were considered by the working group in preparing its final list.

A small subcommittee of the CAGP was organized, with input from representatives from the CAPM and the Canadian Geriatrics Society (CGS). The group reviewed the recommendations made by members of a CPA membership survey, as well as the CGS, AGS and the American Psychiatric Association's (APA) recommendations for *Choosing Wisely*. Two recommendations were selected and discussed, and minor revisions were made to the paragraphs underneath the recommendations. The CAGP also focused the recommendation about benzodiazepines and other hypnotics on insomnia, rather than on a variety of conditions.

The Executive Committee of the Canadian Academy of Child and Adolescent Psychiatry (CACAP) developed a draft list of items after reviewing recommendations made by members of a CPA membership survey, as well as the American Psychiatric Association's (APA) recommendations for *Choosing Wisely*. The list was further discussed and refined and additional feedback was obtained from the CACAP Board of Directors, as well as the Section of Child and Adolescent Psychiatry of the Alberta Psychiatric Association and colleagues elsewhere in the country.

Sources

- 1** Ferracioli-Oda E, Qawasmi A, Bloch MH. Meta-analysis: melatonin for the treatment of primary sleep disorders. *PLoS One*. 2013 May 17;8(5):e63773.
Mindell JA, Owens JA. *A clinical guide to pediatric sleep: Diagnosis and management of sleep problems*. 2nd edition. Philadelphia (PA): Lippincott Williams & Wilkins; 2010.
Morgenthaler TI, Owens J, Alessi C, Boehlecke B, Brown TM, Coleman J, et al. Practice parameters for behavioral treatment of bedtime problems and night wakings in infants and young children: an American Academy of Sleep Medicine report. *Sleep*. 2006;(29)10:1277-81.
Owens JA, Moturi S. Pharmacologic treatment of pediatric insomnia. *Child and Adolescent Psychiatric Clinics of North America*. 2009 Oct;18(4):1001-16.
Stepanski EJ, Wyatt JK. Use of sleep hygiene in the treatment of insomnia. *Sleep Med Rev*. 2003 Jun;7(3):215-25.
- 2** Bhatia SK and Bhatia SC. Childhood and Adolescent depression. *Am Fam Physician*. 2007 Jan 1;75(1):73-80.
Birmaher B, Brent D, AACAP Work Group on Quality Issues, Bernet W, Bukstein O, Walter H, et al. Practice parameter for the assessment and treatment of children and adolescents with depressive disorders. *J Am Acad Child Adolesc Psychiatry*. 2007 Nov;46(11):1503-26.
Hetrick SE, McKenzie JE, Cox GR, Simmons MB, Merry SN. Newer generation antidepressants for depressive disorders in children and adolescents. *Cochrane Database Syst Rev*. 2012 Nov 14;11:CD004851.
Zuckerbrot RA, Cheung AH, Jensen PS, Stein RE, Laraque D; GLAD-PC Steering Group. Guidelines for adolescent depression in primary care (GLAD-PC): 1. Identification, assessment, and initial management. *Pediatrics*. 2007 Nov;120(5): e1299-1312.
- 3** Gorman DA, Gardner DM, Murphy AL, Feldman M, Bélanger SA, Steele MM, et al. Canadian guidelines on pharmacotherapy for disruptive and aggressive behaviour in children and adolescents with attention-deficit hyperactivity disorder, oppositional defiant disorder, or conduct disorder. *Can J Psychiatry*. 2015 Feb;60(2):62-76.
Loy JH, Merry SN, Hetrick SE, Stasiak K. Atypical antipsychotics for disruptive behaviour disorders in children and youths. *Cochrane Database Syst Rev*. 2012 Sep 12;9:CD008559.
Pringsheim T, Hirsch L, Gardner D, Gorman DA. The Pharmacological Management of Oppositional Behaviour, Conduct Problems, and Aggression in Children and Adolescents With Attention-Deficit Hyperactivity Disorder, Oppositional Defiant Disorder, and Conduct Disorder: A Systematic Review and Meta-Analysis. Part 1: Psychostimulants, Alpha-2 Agonists, and Atomoxetine. *Can J Psychiatry*. 2015 Feb 1;60(2):42-51.
Wilkes TCR, Nixon MK. Pharmacological treatment of child and adolescent disruptive behaviour disorders. *Can J Psychiatry*. 2015 Feb;60(2):39-41.
- 4** Canadian ADHD Resource Alliance. Canadian ADHD Practice Guidelines, 3rd Edition [Internet]. 2011 [cited 2015 May 1]. Available from: www.caddra.ca.
Greenhill L, Kollins S, Abikoff H, McCracken J, Riddle M, Swanson J, et al. Efficacy and safety of immediate-release methylphenidate treatment for preschoolers with ADHD. *J Am Acad Child Adolesc Psychiatry*. 2006 Nov;45(11):1284-93.
March JS. The preschool ADHD treatment study (PATS) as the culmination of twenty years of clinical trials in pediatric psychopharmacology. *J Am Acad Child Adolesc Psychiatry*. 2011 May;50(5):427-30.
- 5** Agency for Healthcare Quality and Research. Off-Label Use of Atypical Antipsychotics: An Update [Internet]. 2011 Sep [cited 2015 May 1]. Available from: http://effectivehealthcare.ahrq.gov/ehc/products/150/778/CER43_Off-labelAntipsychotics_20110928.pdf.
Coe HV, Hong IS. Safety of low doses of quetiapine when used for insomnia. *Ann Pharmacother*. 2012 May;46(5):718-22.
Hermes ED, Sernyak M, Rosenheck R. Use of second-generation antipsychotic agents for sleep and sedation: a provider survey. *Sleep*. 2013 Apr;36(4):597-600.
Shah C, Sharma TR, Kablinger A. Controversies in the use of second generation antipsychotics as sleep agent. *Pharmacol Res*. 2014 Jan;79:1-8.

- 6** Korn CS, Currier GW, Henderson SO. "Medical clearance" of psychiatric patients without medical complaints in the emergency department. *J Emerg Med.* 2000 Feb;18(2):173-176.
- Olshaker JS, Browne B, Jerrard DA, Prendergast H, Stair TO. Medical clearance and screening of psychiatric patients in the emergency department. *Acad Emerg Med.* 1997 Feb;4(2):124-128.
- Schiller MJ, Shumway M, Batki SL. Utility of routine drug screening in a psychiatric emergency setting. *Psychiatr Serv.* 2000 Apr;51(4):474-78.
- Tenenbein M. Do you really need that emergency drug screen? *Clin Toxicol.* 2009 Apr;47(4):286-91.
- 7** Barbui C, Cipriani A, Patel V, Ayuso-Mateos JL, van Ommeren M. Efficacy of antidepressants and benzodiazepines in minor depression: systematic review and meta-analysis. *Br J Psychiatry.* 2011 Jan;198(1):11-6.
- Cuijpers P, van Straten A, van Oppen P, Andersson G. Are psychosocial and pharmacologic interventions equally effective in the treatment of adult depressive disorders? A meta-analysis of comparative studies. *J Clin Psychiatry.* 2008 Nov;69(11):1675-85.
- Esposito E, Wang JL, Adair CE, Williams JV, Dobson K, Schopflocher D, et al. Frequency and adequacy of depression treatment in a Canadian population sample. *Can J Psychiatry.* 2007 Dec;52(12):780-789.
- Fournier JC, DeRubeis RJ, Hollon SD, Dimidjian S, Amsterdam JD, Shelton RC, et al. Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA.* 2010 Jan 6;303(1):47-53.
- Kirsch I, Deacon BJ, Huedo-Medina TB, Scoboria A, Moore TJ, Johnson BT. Initial Severity and antidepressant benefits: a meta-analysis of data submitted to the Food and Drug Administration. *PLoS Med.* 2008 Feb;5(2):e45.
- National Institute for Health and Care Excellence. Depression in adults: The treatment and management of depression in adults [Internet]. 2009 Oct [cited 2015 May 1]. Available from: <https://www.nice.org.uk/guidance/cg90>.
- National Institute for Health and Care Excellence. Depression in adults: evidence update [Internet]. 2012 Apr 1 [cited 2015 May 1]. Available from: <https://www.nice.org.uk/guidance/cg90/evidence>.
- 8** Albon E, Tsourapas A, Frew E, Davenport C, Oyeboode F, Bayliss S, et al. Structural neuroimaging in psychosis: a systematic review and economic evaluation. *Health Technol Assess.* 2008 May;12(18):iii-iv, ix-163.
- Goulet K, Deschamps B, Evoy F, Trudel JF. Use of brain imaging (computed tomography and magnetic resonance imaging) in first-episode psychosis: review and retrospective study. *Can J Psychiatry.* 2009 Jul;54(7):493-501.
- Khandanpour N, Hoggard N, Connolly D. The role of MRI and CT of the brain in first episodes of psychosis. *Clin Radiol.* 2013 Mar;68(3):245-50.
- National Institute for Health and Clinical Excellence. Technology appraisal guidance: Structural neuroimaging in first-episode psychosis [Internet]. 2008 Feb [2015 May 1]. Available from: <https://www.nice.org.uk/guidance/ta136>.
- Williams SR, Koyanagi CY, Hishinuma ES. On the usefulness of structural brain imaging for young first episode inpatients with psychosis. *Psychiatry Res.* 2014 Nov 30;224(2):104-6.
- 9** Alessi-Severini S, Bolton JM, Enns MW, Dahl M, Collins DM, Chateau D, et al. Use of benzodiazepines and related drugs in Manitoba: A population-based study. *CMAJ Open.* 2014 Oct;2(4):E208-16.
- Ashton H. The diagnosis and management of benzodiazepine dependence. *Curr Opin Psychiatry.* 2005 May;18(3):249-55.
- Bell CM, Fischer HD, Gill SS, Zagorski B, Sykora K, Wodchis WP, et al. Initiation of benzodiazepines in the elderly after hospitalization. *J Gen Intern Med.* 2007 Jul;22(7):1024-29.
- Cunningham CM, Hanley GM, Morgan S. Patterns in the use of benzodiazepines in British Columbia: Examining the impact on increasing research and guideline cautions against long-term use. *Health Policy.* 2010 Oct;97(2-3):122-9.
- Grad R, Tamblyn R, Holbrook AM, Hurley J, Feightner J, Gayton D. Risk of a new benzodiazepine prescription in relation to recent hospitalization. *J Am Geriatr Soc.* 1999 Feb;47(2):184-8.
- Lader M. Benzodiazepines revisited-will we ever learn? *Addiction.* 2011 Dec;106(12):2086-109.
- Olson M, King M, Schoenbaum M. Benzodiazepine use in the United States. *JAMA Psychiatry.* 2015 Feb;72(2):136-42.
- Olson M, King M, Schoenbaum M. The popularity of benzodiazepines, their advantages, and inadequate pharmacological alternatives—Reply. *JAMA Psychiatry.* 2015 Apr 1.
- Swinson R, Antony M, Katzman MA, Bleay P, Chokka P, Craven M, et al. Clinical practice guidelines: Management of anxiety disorders. *Can J Psychiatry.* 2006 Jul;51 Suppl 2:1S-93S.
- Yokoi Y, Misal M, Oh E, Bellantoni M, Rosenberg PB. Benzodiazepine discontinuation and patient outcome in a chronic geriatric medical/psychiatric unit: a retrospective chart review. *Geriatr Gerontol Int.* 2014 Apr;14(2):388-94.
- 10** Hashimoto E, Tayama M, Ishikawa H, Yamamoto M, Saito T. Influence of comorbid alcohol use disorder on treatment response of depressive patients. *J Neural Transm.* 2015 Feb;122(2):301-6.
- McIntosh C, Ritson B. Treating depression complicated by substance misuse. *Adv Psychiatr Treat.* 2001 Jan;7(5):357-64.
- National Institute for Health and Care Excellence. Alcohol-use disorders: diagnosis, assessment and management of harmful drinking and alcohol dependence [Internet]. 2011 Feb [cited 2015 May 1]. Available from: <http://www.nice.org.uk/guidance/cg115>.
- Nunes EV, Levin FR. Treatment of co-occurring depression and substance dependence: Using meta-analysis to guide clinical recommendations. *Psychiatr Ann.* 2008 Nov;38(11):nihpa128505.
- Petrakis IL, Gonzalez G, Rosenheck R, Krystal JH. Comorbidity of alcoholism and psychiatric disorders. *Alcohol Research and Health.* 2002 Nov;81-9.
- Torrens M, Fonseca F, Mateu G, Farré M. Efficacy of antidepressants in substance use disorders with and without comorbid depression: a systematic review and meta-analysis. *Drug Alcohol Depend.* 2005 Apr 4;78(1):1-22.
- 11** Canadian Agency for Drugs and Technologies in Health. A systematic review of combination and high-dose atypical antipsychotic therapy in patients with schizophrenia. Optimal Use Report: CADTH Volume 1, Issue 1B [Internet]. 2011 Dec [cited 2015 May 1]. Available from: http://www.cadth.ca/media/pdf/H0503_AAP_science-report_e.pdf.
- Fisher MD, Reilly K, Isenberg K, Villa KF. Antipsychotic patterns of use in patients with schizophrenia: polypharmacy versus monotherapy. *BMC Psychiatry.* 2014 Nov 30;14:341.

- 12** Banerjee S. The use of antipsychotic medication for people with dementia: Time for action [Internet]. 2009 Oct [cited 2015 May 1]. Available from: http://webarchive.nationalarchives.gov.uk/20130107105354/http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/documents/digitalasset/dh_108302.pdf.
- Brodaty H, Arasaratnam C. Meta-analysis of nonpharmacological interventions for neuropsychiatric symptoms of dementia. *Am J Psychiatry*. 2012 Sep;169(9):946-53.
- Gill SS, Bronskill SE, Normand SL, Anderson GM, Sykora K, Lam K, et al. Antipsychotic drug use and mortality in older adults with dementia. *Ann Intern Med*. 2007 Jun 5;146(11):775-86.
- Gill SS, Rochon PA, Herrmann N, Lee PE, Sykora K, Gunraj N, et al. Atypical antipsychotic drugs and risk of ischaemic stroke: population based retrospective cohort study. *BMJ*. 2005 Feb 26;330(7489):445.
- Lee PE, Gill SS, Freedman M, Bronskill SE, Hillmer MP, Rochon PA. Atypical antipsychotic drugs in the treatment of behavioural and psychological symptoms of dementia: systematic review. *BMJ*. 2004 Jul 10;329(7457):75.
- Schneider LS, Dagerman K, Insel PS. Efficacy and adverse effects of atypical antipsychotics for dementia: meta-analysis of randomized, placebo-controlled trials. *Am J Geriatr Psychiatry*. 2006 Mar;14(3):191-210.
- Seitz DP, Brisbin S, Herrmann N, Rapoport MJ, Wilson K, Gill SS, et al. Efficacy and feasibility of nonpharmacological interventions for neuropsychiatric symptoms of dementia in long term care: a systematic review. *J Am Med Dir Assoc*. 2012 Jul;13(6):503,506.e2.
- 13** Allain H, Bentué-Ferrer D, Polard E, Akwa Y, Patat A. Postural instability and consequent falls and hip fractures associated with use of hypnotics in the elderly: a comparative review. *Drugs Aging*. 2005;22(9):749-65.
- American Geriatrics Society 2012 Beers Criteria Update Expert Panel. American Geriatrics Society updated Beers Criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc*. 2012 Apr;60(4):616-31.
- Finkle WD, Der JS, Greenland S, Adams JL, Ridgeway G, Blaschke T, et al. Risk of fractures requiring hospitalization after an initial prescription for zolpidem, alprazolam, lorazepam, or diazepam in older adults. *J Am Geriatr Soc*. 2011 Oct;59(10):1883-90.
- Glass J, Lanctôt KL, Herrmann N, Sproule BA, Busto UE. Sedative hypnotics in older people with insomnia: meta-analysis of risks and benefits. *BMJ*. 2005 Nov 19;331(7526):1169.
- McMillan JM, Aitken E, Holroyd-Leduc JM. Management of insomnia and long-term use of sedative-hypnotic drugs in older patients. *CMAJ*. 2013 Nov 19;185(17):1499-505.
- Rapoport MJ, Lanctôt KL, Streiner DL, Bédard M, Vingilis E, Murray B, et al. Benzodiazepines and driving: a meta-analysis. *J Clin Psychiatry*. 2009 Apr 21;70(5):663-673.
- Roszkowska J, Geraci SA. Management of insomnia in the geriatric patient. *Am J Med*. 2010 Dec;123(12):1087-90.

About Choosing Wisely Canada

Choosing Wisely Canada is a campaign to help physicians and patients engage in conversations about unnecessary tests, treatments and procedures, and to help physicians and patients make smart and effective choices to ensure high-quality care.

For more information on *Choosing Wisely Canada* or to see other lists of Five Things Physicians and Patients Should Question, visit www.choosingwiselycanada.org. Join the conversation on Twitter @ChooseWiselyCA.

About The Canadian Psychiatric Association

The Canadian Psychiatric Association (CPA) is a proud partner of the *Choosing Wisely Canada* campaign. CPA is the national voluntary professional association for Canada's 4,700 psychiatrists and 900 residents. As the national voice of Canada's psychiatrists, the CPA advocates for the professional needs of its members in meeting the mental health needs of Canadians, and promotes excellence in education, research and clinical practice. Its mission is to provide a strong, collective voice for psychiatrists across the country and to foster a community dedicated to ensuring the highest possible standards of professional practice in providing psychiatric services to Canadians.

About The Canadian Academy of Child Psychiatry

The Canadian Academy of Child Psychiatry (CACAP) is a proud partner of the *Choosing Wisely Canada* campaign. CACAP promotes quality care and service to the children, youth and families of Canadians within an approach that includes the biological, the psychological and the social; that works with other professional disciplines; and across many sectors of health and other related service organizations.

About The Canadian Academy of Geriatric Psychiatry

The Canadian Academy of Geriatric Psychiatry (CAGP) is a proud partner of the *Choosing Wisely Canada* campaign. CAGP is a national organization of psychiatrists dedicated to promoting mental health in the Canadian elderly population through the clinical, educational, research and advocacy activities of its membership. It was founded in 1991, and is recognized as the voice of Geriatric Psychiatry in Canada. The CAGP is a member of the Council of Academies of the Canadian Psychiatric Association. There are over 300 current members.